

Chapter 13: Complex Numbers

Sections 13.3 & 13.4

1. Function of a complex variable

- A **(single-valued) function** f of a complex variable z is such that for every z in the **domain of definition** \mathcal{D} of f , there is a unique complex number w such that

$$w = f(z).$$

- The **real and imaginary parts** of f , often denoted by u and v , are such that

$$f(z) = u(x, y) + iv(x, y), \quad z = x + iy, \quad x, y \in \mathbb{R}, \quad f(z) \in \mathbb{C},$$

with $u(x, y) \in \mathbb{R}$ and $v(x, y) \in \mathbb{R}$.

Function of a complex variable (continued)

- **Examples:**

- $f(z) = z$ is such that $u(x, y) = x$ and $v(x, y) = y$.
- Find the real and imaginary parts of $f(z) = \bar{z}$.
- $f(z) = \frac{1}{z}$ is defined for all $z \neq 0$ and is such that

$$u(x, y) = \frac{x}{x^2 + y^2}, \quad v(x, y) = \frac{y}{x^2 + y^2}.$$

2. Limits and continuity

- An **open neighborhood** of the point $z_0 \in \mathbb{C}$ is a set of points $z \in \mathbb{C}$ such that

$$|z - z_0| < \epsilon, \quad \text{for some } \epsilon > 0.$$

- Let f be a function of a complex variable z , defined in a neighborhood of $z = z_0$, except maybe at $z = z_0$.
- We say that f has the **limit** w_0 as z goes to z_0 , i.e. that

$$\lim_{z \rightarrow z_0} f(z) = w_0,$$

if for every $\epsilon > 0$, one can find $\delta > 0$, such that for all $z \in \mathcal{D}$,

$$|z - z_0| < \delta \implies |f(z) - w_0| < \epsilon.$$

- **Example:** $\lim_{z \rightarrow i} \frac{z^2 + 1}{z - i} = 2i.$

Continuity

- The function f is **continuous** at $z = z_0$ if f is defined in a neighborhood of z_0 (including at $z = z_0$), and

$$\lim_{z \rightarrow z_0} f(z) = f(z_0).$$

- If $f(z)$ is continuous at $z = z_0$, so is $\overline{f(z)}$. Therefore, if f is continuous at $z = z_0$, so are $\Re(f)$, $\Im(f)$, and $|f|^2$.
- Conversely, if $u(x, y)$ and $v(x, y)$ are continuous at (x_0, y_0) , then $f(z) = u(x, y) + iv(x, y)$ with $z = x + iy$, is continuous at $z_0 = x_0 + iy_0$.
- **Example:** Is the function such that $f(z) = \Im(z^2)/|z|^2$ for $z \neq 0$ and $f(0) = 0$, continuous at $z = 0$?

3. Differentiability

- Assume that f is defined in a neighborhood of $z = z_0$. The **derivative** of the function f at $z = z_0$ is

$$f'(z_0) = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0},$$

assuming that this limit exists.

- If f has a derivative at $z = z_0$, we say that f is **differentiable** at $z = z_0$.
- **Examples:**
 - $f(z) = \bar{z}$ is continuous but not differentiable at $z = 0$.
 - $f(z) = z^3$ is differentiable at any $z \in \mathbb{C}$ and $f'(z) = 3z^2$.

Rules for continuity, limits and differentiation

- To find the limit or derivative of a function $f(z)$, **proceed as you would do for a function of a real variable.**

- **Examples:**

- $f' \left(\frac{1}{z} \right) = -\frac{1}{z^2}.$

- $\frac{d}{dz} z^n = n z^{n-1}, \quad n \in \mathbb{N}.$

- Find $\lim_{z \rightarrow -i} \left(z + \frac{1}{z} \right).$

Rules for continuity, limits and differentiation (continued)

- Properties involving the sum, difference or product of functions of a complex variable are the same as for functions of a real variable. In particular,
 - The limit of a product (sum) is the product (sum) of the limits.
 - The **product and quotient rules** for differentiation still apply.
 - The **chain rule** still applies.

- **Examples:**

- Find $\frac{d}{dz} \left(\frac{z^2 + 1}{z - i} \right).$

- Find $\frac{d}{dz} (z^3 + 9z - 7)^4.$

4. Analytic functions

- A function $f(z)$ is **analytic** at $z = z_0$ if $f(z)$ is differentiable in a neighborhood of z_0 .
- A **region** of the complex plane is a set consisting of an open set, possibly together with some or all of the points on its boundary.
- We say that f is analytic in a region \mathcal{R} of the complex plane, if it is analytic at every point in \mathcal{R} .
- One may use the word **holomorphic** instead of the word analytic.

Analytic functions (continued)

- A function that is analytic at every point in the complex plane is called **entire**.
- **Polynomials** of a complex variable are entire.
 - For instance, $f(z) = 3z - 7z^2 + z^3$ is analytic at every z .
- **Rational functions** of a complex variable of the form $f(z) = \frac{g(z)}{h(z)}$, where g and h are polynomials, are analytic everywhere, except at the zeros of $h(z)$.
 - For instance, $\frac{z^2 + 1}{z - i}$ is analytic except at $z = i$.
 - In the above example, $z = i$ is called a **pole** of $f(z)$.

5. The Cauchy-Riemann equations

- If $f(z) = u(x, y) + iv(x, y)$ is defined in a neighborhood of $z = x + iy$, and if f is differentiable at z , then

$$u_x(x, y) = v_y(x, y), \quad \text{and} \quad u_y(x, y) = -v_x(x, y). \quad (1)$$

These are called **the Cauchy-Riemann equations**.

- Conversely, if the partial derivatives of u and v exist in a neighborhood of $z = x + iy$, if they are continuous at z and satisfy the Cauchy-Riemann equations at z , then $f(z) = u(x, y) + iv(x, y)$ is differentiable at z .
- The Cauchy-Riemann equations therefore give a **criterion for analyticity**.

The Cauchy-Riemann equations (continued)

- Indeed, if a function is analytic at z , it must satisfy the Cauchy-Riemann equations in a neighborhood of z . In particular, **if f does not satisfy the Cauchy-Riemann equations, then f cannot be analytic**.
- Conversely, if the partial derivatives of u and v exist, are continuous, and satisfy the Cauchy-Riemann equations in a neighborhood of $z = x + iy$, then $f(z) = u(x, y) + iv(x, y)$ is analytic at z .
- **Examples:**
 - Use the Cauchy-Riemann equations to show that \bar{z} is not analytic.
 - Use the Cauchy-Riemann equations to show that $\frac{1}{z}$ is analytic everywhere except at $z = 0$.

Applications of the Cauchy-Riemann equations

- A consequence of the Cauchy-Riemann equations is that

$$f'(z) = u_x + iv_x = v_y - iu_y. \quad (2)$$

- We will use these formulas later to calculate the derivative of some analytic functions.
- Another consequence of the Cauchy-Riemann equations is that an entire function with constant absolute value is constant. In fact, a more general result is that **an entire function that is bounded** (including at infinity) **is constant**.

6. Harmonic functions

- One can show that if f is analytic in a region \mathcal{R} of the complex plane, then it is infinitely differentiable at any point in \mathcal{R} .
- If $f(z) = u(x, y) + iv(x, y)$ is analytic in \mathcal{R} , then both u and v satisfy **Laplace's equation** in \mathcal{R} , i.e.

$$\nabla^2 u = u_{xx} + u_{yy} = 0, \quad \text{and} \quad \nabla^2 v = v_{xx} + v_{yy} = 0. \quad (3)$$

- A function that satisfies Laplace's equation is called an **harmonic function**.

Harmonic conjugate

- If $f(z) = u(x, y) + iv(x, y)$ is analytic in \mathcal{R} , then we saw that both u and v are harmonic (i.e. satisfy Laplace's equation) in \mathcal{R} .
- We say that u and v are **harmonic conjugates** of one another.
- Given an harmonic function u , one can use the Cauchy-Riemann equations to find its harmonic conjugate v , and vice-versa.
- **Examples:**
 - Check that $u(x, y) = 2xy$ is harmonic, and find its harmonic conjugate v .
 - Given an harmonic function $v(x, y)$, how would you find its harmonic conjugate $u(x, y)$?